Translated English of Chinese Standard: GB31604.30-2016

www.ChineseStandard.net → Buy True-PDF → Auto-delivery.

Sales@ChineseStandard.net

GB

NATIONAL STANDARD OF THE PEOPLE'S REPUBLIC OF CHINA

GB 31604.30-2016

National Food Safety Standard – Food Contact

Materials and Articles – Determination of the Content

and Migration of Phthalate Esters

Issued on: October 19, 2016 Implemented on: April 19, 2017

Issued by: National Health and Family Planning Commission of PRC

Table of Contents

reword	4
Scope	5
Principle	5
Reagents and Materials	5
Apparatus	6
Analytical Procedures	7
Expression of Analytical Results	8
Precision	9
Others	9
Principle	9
Reagents and Materials	10
Apparatus	10
Analytical Procedures	11
Expression of Analytical Results	12
Precision	13
Others	13
Appendix A Information on 18 kinds of Commonly Used Phthalate Ester	
sticizers	14
ppendix B Quantitative and Qualitative Selective Ion List of Phthalate Est	er
mpounds	15
Appendix C Gas Chromatography – Mass Spectrometry Total Ion	
romatogram of Phthalate Ester Standard Sample Solution	16
	Scope Principle Reagents and Materials Apparatus Analytical Procedures Expression of Analytical Results Precision Others Principle Reagents and Materials Apparatus Analytical Procedures Expression of Analytical Results Apparatus Apparatus Apparatus Apparatus Expression of Analytical Results Precision Others Appendix A Information on 18 kinds of Commonly Used Phthalate Ester sticizers Appendix B Quantitative and Qualitative Selective Ion List of Phthalate Ester prounds Appendix C Gas Chromatography – Mass Spectrometry Total Ion

National Food Safety Standard – Food Contact Materials and Articles – Determination of the Content and Migration of Phthalate Esters

1 Scope

This Standard specifies the determination of the content and migration of phthalate esters in the food contact materials and articles.

This Standard is applicable to the determination of the content of phthalate esters in the food plastic packaging materials, and the determination of migration of phthalate esters in food contact materials and articles.

Determination of Phthalate Esters

2 Principle

After crushing the food plastic packaging materials and articles, use n-hexane to conduct the ultrasonic extraction; after filtering the extracting solution, use gas chromatography-mass spectrometry to determine. Adopt the feature selective ion monitoring scan mode (SIM), so that retain the qualitative of abundance ratio of the time and fragments; use external standard method to quantify.

3 Reagents and Materials

Unless otherwise is specified, the reagents used in this Method are all chromatographic pure (or analytically pure that has been redistilled). Water shall be the one redistilled by a full glass device and stored in the glass vessel.

3.1 Reagents

- **3.1.1** N-hexane (C_6H_{14}) .
- 3.12 Acetone (CH₃COCH₃).

3.2 Standard substance of phthalate esters

5 Analytical Procedures

5.1 Specimen preparation

Take 5g of typical sample; shear the specimen (excluding the preservative film) into single pieces with diameter≤0.2cm; the preservative film piece diameter≤0.3cm; then mix evenly; accurately take 0.2g~0.5g of specimen (accurate to 0.0001g) into stoppered triangular flask; add 20mL of n-hexane; perform ultrasonic extraction for 30min and filter; after than the residue is re-extracted once by 20mL of n-hexane; combine the filtrate into 50mL volumetric flask; use n-hexane to make constant volume to the scale; then dilute it as per the content of phthalate esters in the specimen; after mixing evenly, pass through a 0.45µm organic phase glass filter film; and be analyzed in the gas chromatograph-mass spectrometer.

5.2 Blank test

The reagents used in the test shall be treated as per 5.1; and analyzed in GC-MS.

5.3 Instrument reference conditions

5.3.1 Gas chromatography reference conditions

- a) Chromatographic column: 5% polyphenylmethylsiloxane quartz capillary column or analytical column with similar performance; the specification indicates column length of 30.0m, inner diameter of 0.25mm, film thickness of 0.25µm;
- b) Sample inlet temperature: 260°C;
- c) Temperature rise program: maintain the initial column temperature of 60°C for 1min; raise temperature to 220°C by 20°C/min, maintain for 1min; raise temperature to 250°C by 5°C/min, maintain for 1min; then raise temperature to 290°C by 20°C/min, maintain for 7.5min;
- d) Carrier gas: helium (purity >99.999%), flow rate: 1mL/min;
- e) Sampling injecting method: splitless sample injection;
- f) Sample injecting volume: 1µL.

5.3.2 Mass spectrometry reference conditions

- a) Gas chromatography and mass spectrometry interface temperature: 280°C;
- b) Ion source temperature: 230°C;
- c) Ionization mode: electron impact ionization source (EI);

10 Reagents and Materials

Unless otherwise is specified, the reagents used in this Method are all chromatographic pure (or analytically pure that has been redistilled). Water shall be the one redistilled by a full glass device and stored in the glass vessel.

10.1 Reagents

- **10.1.1** N-hexane (C_6H_{14}) .
- **10.1.2** Absolute alcohol (C₂H₅OH).
- 10.1.3 Acetic acid (CH₃COOH).
- **10.1.4** Reagents required for preparing the acidic and alcoholic food simulants: abide by the provisions of GB 31604.1.
- **10.1.5** Isooctane: oil-based food simulants.

10.2 Standard substance

The standard substance for 18 kinds of phthalate esters, the same as 3.2.

10.3 Preparation of standard solution

- **10.3.1** Standard stock solution: the same as 3.3.1.
- **10.3.2** 17 kinds of phthalate esters standard use solutions: the same as 3.3.2.
- **10.3.3** Diisononyl ortho-phthalate standard use solution: the same as 3.3.3.
- **10.3.4** Phthalate ester standard serial working solution: accurately pipette appropriate amount of phthalate ester standard use solution; use n-hexane to prepare 17 kinds of phthalate ester mixing standard use solutions with concentrations of 0.1mg/L, 0.2mg/L, 0.5mg/L, 1.0mg/L, 3.0mg/L, 8.0mg/L, 10.0mg/L, as well as single-labeled diisononyl ortho-phthalate (DINP) with concentrations of 0.5mg/L, 1.0mg/L, 3.0mg/L, 8.0mg/L, 10.0mg/L for later-use.

11 Apparatus

- **11.1** Gas chromatography-mass spectrometer: with electron impact source (EI).
- 11.2 Centrifuge.
- 11.3 Oscillator.

- **11.4** Solid phase extraction device.
- **11.5** Constant temperature oven.
- **11.6** Low-temperature incubator.
- 11.7 Centrifuge tube: 10mL.
- 11.8 Volumetric flask: 10mL.
- 11.9 Glassware.

NOTE: after washing the used glassware, rinse by the distilled water for 3 times; soak in acetone for 1h; bake at 200°C for 2h; cool off to the room temperature for later-use.

12 Analytical Procedures

12.1 Preparation of food simulant test solution

According to the requirements of GB 5009.156 and GB 31604.1, perform the migration test against the sample; then obtain the food simulant test solution.

12.2 Treatment of soaking solution

Place the soaking solution at the room temperature; for the isooctane and 95% ethanol aqueous solution (≥ 50% alcoholic food) simulant soaking solution, accurately take 10g±0.01g of soaking solution into the pear-shaped distillation flask; reduce pressure in 45°C water bath; perform rotary evaporation to dryness; add 1mL of n-hexane to dissolve; perform vortex oscillation for 3min; centrifuge for 5min at 4000r/min; collect the supernatant for the instrument test.

For other simulant soaking solutions (including water-based, acidic food, <50% alcoholic food simulant), take 10g±0.01g of simulant soaking solution into 25mL centrifuge tube; add 4mL of n-hexane; shake and extract for 10min; centrifuge for 5min at 4000r/min speed; take out the upper layer of n-hexane; repeatedly extract twice; combine the n-hexane phase into pear-shaped distillation flask; reduce pressure in 45°C water bath; perform rotary evaporation to dryness; add 1mL of n-hexane to dissolve; perform vortex oscillation for 3min; centrifuge for 5min at 4000r/min; collect the supernatant for the instrument test.

The sample solution can be diluted as per the specific conditions, so that the measured value shall be within the linear range of the standard curve.

12.3 Blank test

The reagents used in the test shall be treated as per 12.2; perform the GC-MS analysis.

This is an excerpt of the PDF (Some pages are marked off intentionally)

Full-copy PDF can be purchased from 1 of 2 websites:

1. https://www.ChineseStandard.us

- SEARCH the standard ID, such as GB 4943.1-2022.
- Select your country (currency), for example: USA (USD); Germany (Euro).
- Full-copy of PDF (text-editable, true-PDF) can be downloaded in 9 seconds.
- Tax invoice can be downloaded in 9 seconds.
- Receiving emails in 9 seconds (with download links).

2. https://www.ChineseStandard.net

- SEARCH the standard ID, such as GB 4943.1-2022.
- Add to cart. Only accept USD (other currencies https://www.ChineseStandard.us).
- Full-copy of PDF (text-editable, true-PDF) can be downloaded in 9 seconds.
- Receiving emails in 9 seconds (with PDFs attached, invoice and download links).

Translated by: Field Test Asia Pte. Ltd. (Incorporated & taxed in Singapore. Tax ID: 201302277C)

About Us (Goodwill, Policies, Fair Trading...): https://www.chinesestandard.net/AboutUs.aspx

Contact: Wayne Zheng, Sales@ChineseStandard.net

Linkin: https://www.linkedin.com/in/waynezhengwenrui/

---- The End -----